

# CHAPTER - 9

## COORDINATION COMPOUNDS - INTEXT SOLUTIONS

### Question 9.1:

Write the formulas for the following coordination compounds:

- (i) Tetraamminediaquacobalt(III) chloride
- (ii) Potassium tetracyanonickelate(II)
- (iii) Tris(ethane-1,2-diamine) chromium(III) chloride
- (iv) Amminebromidochloridonitrito-N-platinate(II)
- (v) Dichloridobis(ethane-1,2-diamine)platinum(IV) nitrate
- (vi) Iron(III) hexacyanoferrate(II)

Answer

- (i)  $[\text{Co}(\text{H}_2\text{O})_2(\text{NH}_3)_4]\text{Cl}_3$
- (ii)  $\text{K}_2[\text{Ni}(\text{CN})_4]$
- (iii)  $[\text{Cr}(\text{en})_3]\text{Cl}_3$
- (iv)  $[\text{Pt}(\text{NH}_3)_3\text{BrCl}(\text{NO}_2)]^-$
- (v)  $[\text{PtCl}_2(\text{en})_2](\text{NO}_3)_2$
- (vi)  $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$

### Question 9.2:

Write the IUPAC names of the following coordination compounds:

- (i)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$
- (ii)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
- (iii)  $\text{K}_3[\text{Fe}(\text{CN})_6]$
- (iv)  $\text{K}_3[\text{Fe}(\text{C}_2\text{O}_4)_3]$
- (v)  $\text{K}_2[\text{PdCl}_4]$
- (vi)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}(\text{NH}_2\text{CH}_3)]\text{Cl}$

Answer


- (i) Hexaamminecobalt(III) chloride
  - (ii) Pentaamminechloridocobalt(III) chloride
  - (iii) Potassium hexacyanoferrate(III)
  - (iv) Potassium trioxalatoferrate(III)
  - (v) Potassium tetrachloridopalladate(II)
  - (vi) Diamminechlorido(methylamine)platinum(II) chloride
- Question 9.3:**

Indicate the types of isomerism exhibited by the following complexes and draw the structures for these isomers:


- i.  $K[Cr(H_2O)_2(C_2O_4)_2]$
- ii.  $[Co(en)_3]Cl_3$
- iii.  $[Co(NH_3)_5(NO_2)](NO_3)_2$
- iv.  $[Pt(NH_3)(H_2O)Cl_2]$

Answer


- i. Both geometrical (*cis*-, *trans*-) isomers for  $K[Cr(H_2O)_2(C_2O_4)_2]$  can exist. Also, optical isomers for *cis*-isomer exist.



*Trans*-isomer is optically inactive. On the other hand, *cis*-isomer is optically active.




- (ii) Two optical isomers for  $[Co(en)_3]Cl_3$




exist.


Two optical isomers are possible for this structure.



A pair of optical isomers:




It can also show linkage isomerism.



It can also show ionization isomerism.



(iv) Geometrical (*cis*-, *trans*-) isomers of  $[\text{Pt}(\text{NH}_3)(\text{H}_2\text{O})\text{Cl}_2]$  can exist.







#### Question 9.4:

Give evidence that  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{SO}_4$  and  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Cl}$  are ionization isomers.

Answer

When ionization isomers are dissolved in water, they ionize to give different ions. These ions then react differently with different reagents to give different products.




**Question 9.5:**


Explain on the basis of valence bond theory that  $[\text{Ni}(\text{CN})_4]^{2-}$  ion with square planar structure is diamagnetic and the  $[\text{NiCl}_4]^{2-}$  ion with tetrahedral geometry is paramagnetic.

Answer


Ni is in the +2 oxidation state i.e., in  $d^8$  configuration.



There are 4  $\text{CN}^-$  ions. Thus, it can either have a tetrahedral geometry or square planar geometry. Since  $\text{CN}^-$  ion is a strong field ligand, it causes the pairing of unpaired  $3d$  electrons.



It now undergoes  $dsp^2$  hybridization. Since all electrons are paired, it is diamagnetic. In case of  $[\text{NiCl}_4]^{2-}$ ,  $\text{Cl}^-$  ion is a weak field ligand. Therefore, it does not lead to the pairing of unpaired  $3d$  electrons. Therefore, it undergoes  $sp^3$  hybridization.




Since there are 2 unpaired electrons in this case, it is paramagnetic in nature.

**Question 9.6:**


$[\text{NiCl}_4]^{2-}$  is paramagnetic while  $[\text{Ni}(\text{CO})_4]$  is diamagnetic though both are tetrahedral. Why?

Answer

Though both  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CO})_4]$  are tetrahedral, their magnetic characters are different. This is due to a difference in the nature of ligands.  $\text{Cl}^-$  is a weak field ligand and it does not cause the pairing of unpaired  $3d$  electrons. Hence,  $[\text{NiCl}_4]^{2-}$  is paramagnetic.



In  $[\text{Ni}(\text{CO})_4]$ , Ni is in the zero oxidation state i.e., it has a configuration of  $3d^8 4s^2$ .



But CO is a strong field ligand. Therefore, it causes the pairing of unpaired  $3d$  electrons. Also, it causes the  $4s$  electrons to shift to the  $3d$  orbital, thereby giving rise to  $sp^3$  hybridization. Since no unpaired electrons are present in this case,  $[\text{Ni}(\text{CO})_4]$  is diamagnetic.


**Question 9.7:**

$[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$  is strongly paramagnetic whereas  $[\text{Fe}(\text{CN})_6]^{3-}$  is weakly paramagnetic.

Explain.

Answer

In both  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$  and  $[\text{Fe}(\text{CN})_6]^{3-}$ , Fe exists in the +3 oxidation state i.e., in  $d^5$  configuration.



Since  $\text{CN}^-$  is a strong field ligand, it causes the pairing of unpaired electrons. Therefore, there is only one unpaired electron left in the  $d$ -orbital.



Therefore,

$$\begin{aligned}\mu &= \sqrt{n(n+2)} \\ &= \sqrt{1(1+2)} \\ &= \sqrt{3} \\ &= 1.732 \text{ BM}\end{aligned}$$

On the other hand, H<sub>2</sub>O is a weak field ligand. Therefore, it cannot cause the pairing of electrons. This means that the number of unpaired electrons is 5.

Therefore,

$$\begin{aligned}\mu &= \sqrt{n(n+2)} \\ &= \sqrt{5(5+2)} \\ &= \sqrt{35} \\ &\approx 6 \text{ BM}\end{aligned}$$





Thus, it is evident that  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$  is strongly paramagnetic, while  $[\text{Fe}(\text{CN})_6]^{3-}$  is weakly paramagnetic.

**Question 9.8:**

Explain  $[\text{Co}(\text{NH}_3)_6]^{3+}$  is an inner orbital complex whereas  $[\text{Ni}(\text{NH}_3)_6]^{2+}$  is an outer orbital complex.

Answer


|  |  |
|--|--|
| $[\text{Co}(\text{NH}_3)_6]^{3+}$          | $[\text{Ni}(\text{NH}_3)_6]^{2+}$          |
| Oxidation state of cobalt = +3             | Oxidation state of Ni = +2                 |
| Electronic configuration of cobalt = $d^6$ | Electronic configuration of nickel = $d^8$ |

|   |  |
|---|--|
|  <p>NH<sub>3</sub> being a strong field ligand causes the pairing. Therefore, Ni can undergo <math>d^2sp^3</math> hybridization.</p>  <p>Hence, it is an inner orbital complex.</p> |  <p>If NH<sub>3</sub> causes the pairing, then only one 3d orbital is empty. Thus, it cannot undergo <math>d^2sp^3</math> hybridization. Therefore, it undergoes <math>sp^3d^2</math> hybridization.</p>  <p>Hence, it forms an outer orbital complex.</p> |
|---|--|


**Question 9.9:**

Predict the number of unpaired electrons in the square planar  $[\text{Pt}(\text{CN})_4]^{2-}$  ion.

Answer



In this complex, Pt is in the +2 state. It forms a square planar structure. This means that it undergoes  $dsp^2$  hybridization. Now, the electronic configuration of  $\text{Pt}(+2)$  is  $5d^8$ .





$\text{CN}^-$  being a strong field ligand causes the pairing of unpaired electrons. Hence, there are

no unpaired electrons in  $[\text{Pt}(\text{CN})_4]^{2-}$ .

**Question 9.10:**

The hexaquo manganese(II) ion contains five unpaired electrons, while the hexacyanoion contains only one unpaired electron. Explain using Crystal Field Theory. Answer



Mn is in the +2 oxidation state.

Mn is in the +2 oxidation state.

The electronic configuration is  $d^5$ .

The electronic configuration is  $d^5$ .

The crystal field is octahedral. Water is a weak field ligand. Therefore, the arrangement of the electrons in

The crystal field is octahedral. Cyanide is

arrangement of the electrons in

$[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$  is  $t_{2g}^3 e_g^2$ .

$[\text{Mn}(\text{CN})_6]^{4-}$  is

$t_{2g}^5 e_g^0$ .

Hence, hexaquo manganese (II) ion has five unpaired electrons, while hexacyano ion has only one unpaired electron.

#### Question 9.11:

Calculate the overall complex dissociation equilibrium constant for the  $\text{Cu}(\text{NH}_3)_4^{2+}$  ion, given that  $\beta_4$  for this complex is  $2.1 \times 10^{13}$ . Answer

$$\beta_4 = 2.1 \times 10^{13}$$

The overall complex dissociation equilibrium constant is the reciprocal of the overall stability constant,  $\beta_4$ .

$$\begin{aligned} \frac{1}{\beta_4} &= \frac{1}{2.1 \times 10^{13}} \\ \therefore &= 4.7 \times 10^{-14} \end{aligned}$$